

Vestibular/Concussion/Imbalance Intake Form

Name: _____ Date: _____

Do you exercise? If so, please describe: _____

Primary Concern:

Date of Onset: _____

Describe the problem that brings you to therapy: _____

History of falls? No Yes If yes how often? _____ When was last fall/what happened? _____

How often do symptoms occur? Daily Weekly Constantly

How long do symptoms last? Seconds Minutes Hours Days

Since then, has your problem: Worsened Improved Same


Have you experienced a recent trauma? No Yes If yes describe _____

Have you been ill recently (cold, flu, ear infection)? No Yes If yes, please describe: _____

Have you ever experienced this problem before? No Yes If yes, please describe: _____

Pain/Dizziness (circle one): if you have no pain, please mark as 0.

If you are having pain, please rate the severity on a 0-10 scale, where 0 is no pain and 10 is the most severe pain:											
At Worst	0	1	2	3	4	5	6	7	8	9	10
At Best	0	1	2	3	4	5	6	7	8	9	10
Current	0	1	2	3	4	5	6	7	8	9	10



Symptoms:

Symptom Description (circle all that apply):

Spinning	<input type="checkbox"/>	Rocking/Swaying	<input type="checkbox"/>	Nausea/Vomiting	<input type="checkbox"/>	Balance difficulty	<input type="checkbox"/>
Light Headedness	<input type="checkbox"/>	Ringing in ears	<input type="checkbox"/>	Hearing loss	<input type="checkbox"/>	Ear fullness/ pressure	<input type="checkbox"/>
Headaches	<input type="checkbox"/>	Decreased concentration	<input type="checkbox"/>	Short term memory loss	<input type="checkbox"/>	Long term memory loss	<input type="checkbox"/>
Light sensitivity	<input type="checkbox"/>	Double vision	<input type="checkbox"/>	Objects appear distorted	<input type="checkbox"/>	Sensitivity to motion/vision	<input type="checkbox"/>
Disorientation	<input type="checkbox"/>	Neck pain	<input type="checkbox"/>	Back pain	<input type="checkbox"/>	Facial numbness	<input type="checkbox"/>
Passing out/ fainting	<input type="checkbox"/>	Fatigue	<input type="checkbox"/>	Weakness	<input type="checkbox"/>	Other:	<input type="checkbox"/>

Symptoms increase with (circle all that apply):

Sit to stand	<input type="checkbox"/>	Turn head	<input type="checkbox"/>	Walking	<input type="checkbox"/>	Bending/ Squatting	<input type="checkbox"/>	Reading	<input type="checkbox"/>	Rolling in bed	<input type="checkbox"/>
Lying to sit	<input type="checkbox"/>	Look up/down	<input type="checkbox"/>	Crowds	<input type="checkbox"/>	Cough/ Sneeze	<input type="checkbox"/>	Driving	<input type="checkbox"/>	Bearing down/ straining	<input type="checkbox"/>

Other (please list): _____

Patient/Guardian Signature

Date

Dizziness Handicap Inventory

Instructions: The purpose of this scale is to identify difficulties that you may be experiencing because of your dizziness. Please check “always”, or “no” or “sometimes” to each question. Answer each question only as it pertains to your dizziness problem.

	Questions	Always	Sometimes	No
P1	Does looking up increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E2	Because of your problem, do you feel frustrated?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F3	Because of your problem, do you restrict your travel for business or pleasure?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
P4	Does walking down the aisle of a supermarket increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F5	Because of your problem, do you have difficulty getting into or out of bed?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F6	Does your problem significantly restrict your participation in social activities, such as going out to dinner, going to movies, dancing or to parties?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F7	Because of your problem, do you have difficulty reading?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F8	Does performing more ambitious activities like sports, dancing, and household chores, such as sweeping or putting dishes away; increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E9	Because of your problem, are you afraid to leave your home without having someone accompany you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E10	Because of your problem, have you been embarrassed in front of others?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
P11	Do quick movements of your head increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F12	Because of your problem, do you avoid heights?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
P13	Does turning over in bed increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F14	Because of your problem, is it difficult for you to do strenuous housework or yard work?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E15	Because of your problem, are you afraid people may think that you are intoxicated?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F16	Because of your problem, is it difficult for you to go for a walk by yourself?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
P17	Does walking down a sidewalk increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E18	Because of your problem, is it difficult for you to concentrate?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F19	Because of your problem, is it difficult for you to walk around your house in the dark?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E20	Because of your problem, are you afraid to stay home alone?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E21	Because of your problem, do you feel handicapped?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E22	Has your problem placed stress on your relationship with members of your family or friends?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E23	Because of your problem, are you depressed?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
F24	Does your problem interfere with your job or household responsibilities?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
P25	Does bending over increase your problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Scoring for Dizziness Handicap Inventory

Eval	Total Functional	Total Emotional	Total Physical	TOTAL SCORE
Reassess #1				
Reassess #2				
Reassess #3				
Reassess #4				

Always = 4

Sometimes = 2

No = 0

P = physical

E = emotional

F = functional

Subscales



Notes:

1. Subjective measure of the patient's perception of handicap due to the dizziness
2. Top score is 100 (maximum perceived disability)
3. Bottom score is 0 (no perceived disability)
4. The following 5 items can be useful in predicting BPPV
 - Does looking up increase your problem?
 - Because of your problem, do you have difficulty getting into or out of bed?
 - Do quick movements of your head increase your problem?
 - Does bending over increase your problem?
5. Can use subscale scores to track change as well

Name: _____ Age/DOB: _____ Date of Injury: _____

Post Concussion Symptom Scale - 0 1 2 3 4 5 6
 No symptoms "0" ----- Moderate "3" ----- Severe "6"

BASELINE (prior to concussion)

Post Concussion: Date: _____

Headache	0 1 2 3 4 5 6	Headache	0 1 2 3 4 5 6
Nausea	0 1 2 3 4 5 6	Nausea	0 1 2 3 4 5 6
Vomiting	0 1 2 3 4 5 6	Vomiting	0 1 2 3 4 5 6
Balance Problems	0 1 2 3 4 5 6	Balance Problems	0 1 2 3 4 5 6
Dizziness	0 1 2 3 4 5 6	Dizziness	0 1 2 3 4 5 6
Fatigue	0 1 2 3 4 5 6	Fatigue	0 1 2 3 4 5 6
Trouble falling asleep	0 1 2 3 4 5 6	Trouble falling asleep	0 1 2 3 4 5 6
Excessive sleep	0 1 2 3 4 5 6	Excessive sleep	0 1 2 3 4 5 6
Loss of sleep	0 1 2 3 4 5 6	Loss of sleep	0 1 2 3 4 5 6
Drowsiness	0 1 2 3 4 5 6	Drowsiness	0 1 2 3 4 5 6
Light sensitivity	0 1 2 3 4 5 6	Light sensitivity	0 1 2 3 4 5 6
Noise sensitivity	0 1 2 3 4 5 6	Noise sensitivity	0 1 2 3 4 5 6
Irritability	0 1 2 3 4 5 6	irritability	0 1 2 3 4 5 6
Sadness	0 1 2 3 4 5 6	Sadness	0 1 2 3 4 5 6
Nervousness	0 1 2 3 4 5 6	Nervousness	0 1 2 3 4 5 6
More emotional	0 1 2 3 4 5 6	More emotional	0 1 2 3 4 5 6
Numbness	0 1 2 3 4 5 6	Numbness	0 1 2 3 4 5 6
Feeling "slow"	0 1 2 3 4 5 6	Feeling "slow"	0 1 2 3 4 5 6
Feeling "foggy"	0 1 2 3 4 5 6	Feeling "foggy"	0 1 2 3 4 5 6
Visual problems	0 1 2 3 4 5 6	Visual problems	0 1 2 3 4 5 6
Difficulty concentrating	0 1 2 3 4 5 6	Difficulty concentrating	0 1 2 3 4 5 6
Difficulty remembering	0 1 2 3 4 5 6	Difficulty remembering	0 1 2 3 4 5 6

TOTAL SCORE _____

TOTAL SCORE _____

Use of the Post-Concussion Symptom Scale: The athlete should fill out the form, on his or her own, in order to give a subjective value for each symptom. This form can be used with each encounter to track the athlete's progress towards the resolution of symptoms. Many athletes may have some of these reported symptoms at a baseline, such as concentration difficulties in the patient with attention-deficit disorder or sadness in an athlete with underlying depression, and must be taken into consideration when interpreting the score. Athletes do not have to be at a total score of zero to return to play if they already have had some symptoms prior to their concussion.